

1. Sexual Orientation
Heterosexual
MSM (men who have sex with men)
Lesbian
Bisexual
Transgender – Refer to Level 3

2. Last Sexual Contact

< 2 Weeks

Continue with screen
Advise repeat
Chlamydia/ Gonorrhoea
test – 2 weeks
HIV/Syphilis test – 3
months

> 2 Weeks

Continue with screen
Repeat HIV/Syphilis -3
months

3. Number of Partners in last 3 months

5 or more refer to
Sexual Health
Promotion for
Brief Interventions
Continue with
screen

Less than 5

Continue with
screen

4. Blood Borne Virus Assessment

IVDU
YES → Refer to Level 3 services
NO → Continue with screen

MSM/ BS
YES → Unprotected anal sex with casual partner in the last year → YES → Refer to Level 3 Services
NO → Continue with screen

CSW
YES → Refer to Level 3 Services
NO → Continue with screen

Endemic Areas
YES → Refer to Level 3 Services
NO → Continue with screen

Continue with screen

5. Check for any contraceptive need

Sampling Methods

Females: Self-taken vulvovaginal swabs
Chlamydia/Gonorrhoea (CHL/GC)
Venous sample for HIV/Syphilis

Lesbian / Bisexual Women: Additional pharyngeal swab for CHL/GC

Males: First void urine sample (last passed urine >1 hour ago)
Venous sample for HIV/SGS/Hep B & C as appropriate
Rectal & Pharyngeal swabs for CHL/GC for MSM